
ENDOVENOUS THERMAL ABLATION OF THE SAPHENOUS VEIN

Patient Instructions

These instructions are to prepare you for the endovenous thermal ablation procedures used to treat incompetent saphenous and varicose veins. The saphenous veins frequently do not function properly in patients with chronic leg pain / swelling, and those with varicose veins. These incompetent saphenous veins can be percutaneously destroyed or ablated with thermal energy without stripping the veins from the legs. Ablation of diseased veins results in immediate improvement of venous circulation in the leg, disappearance of the varicose veins, and relief of leg discomfort / swelling associated with venous hypertension.

The following instructions are to help you prepare for your procedure and speed your recovery.

Before and During the Procedure:

- Please drink *plenty* of water the evening before the procedure so that you will be well-hydrated for the procedure. *Do not eat or drink anything after 12:00 midnight before your procedure.*
- Shave your legs (men generally use a clipper, not a razor) the night before your procedure. On the morning of your procedure, shower and wash your legs thoroughly using hibiclens.
- Take your usual medications on the morning of your procedure with a small sip of water (unless Dr. Hayes has asked you to temporarily stop some of your medications)
- Unlike with most procedures, blood-thinning medications are typically best continued during this minimally-invasive procedure. If you are on blood-thinning medications and have any questions about them, please ask Dr. Hayes.
- Please bring a **complete** list of medications with you on the day of your procedure. It is especially important to notify Dr. Hayes and his staff about any allergies or recent changes in your medications.
- Please wear shorts or loose-fitting clothing to the Vein Center. Also wear slip-on shoes or house slippers. Your foot and leg will be wrapped in ace bandages, and regular shoes and pants will not fit over the post-op dressings.
- You will be sedated during the procedure and should not drive home. Please try and find a responsible adult to drive you to and from the Vein Center.
- Your venous ablation will be performed in one of the Vein Center procedure rooms. You will have monitors attached to record your blood pressure, pulse rate, and oxygen levels. Dr. Hayes and his staff will monitor you at all times. You will be given intravenous conscious sedation, and you may have trouble remembering the procedure.
- Endovenous Thermal ablation is performed using local anesthesia along the course of the diseased vein. There may be some discomfort associated with the placement of the anesthetic, and also with use of the ablation fiber/catheter. This discomfort is generally mild and you will receive additional sedation during these times to lessen any discomfort.
- A surgical incision is generally not required. Only small punctures are made to both place the ablation fiber/catheter and to remove the ropey varicose veins. Sometimes a small incision is made at groin level to expose and ligate a large saphenous vein before positioning the ablation fiber/catheter.
- In order to remove the unsightly and painful varicose veins or other visible veins in the leg, most endovenous ablation procedures are accompanied by micro-phlebectomy, sclerotherapy, or both.
- Ultrasound is necessary during the procedure to access the diseased vein, to insert the ablation fiber/catheter, to place the local anesthetic around the vein, and to position the ablation fiber/catheter perfectly within the termination of the diseased vein.
- The ablation fiber/catheter is inserted inside the diseased vein and positioned to begin the ablation. Energy is delivered to the fiber/ catheter, which is slowly withdrawn through the vein. The heat gently destroys the lining of the vein. There is no pulling, stripping, or avulsion of branches.
- The small wounds are closed with absorbable sutures or steri-strips, sterile dressings are applied, and your leg is wrapped with Ace wraps.

After You're Home:

- Leave the ace wraps on your foot and leg until your ultrasound (two to three days after your procedure). Significant discomfort (especially behind the knee) can mean the ace wraps are making a blister. If the ace wraps feel too tight or pinch you behind the knee, loosen them a little bit. Be sure to put them back on right away.
- You may walk around the house and out of doors as early as the afternoon of the procedure. Please avoid sitting with the feet down or standing still for prolonged intervals. We like you to walk around quite a bit, as this speeds recovery and keeps you from becoming too "stiff". While sitting or lying down, elevate your leg on one or more pillows.
- When the ace wraps are removed after your post-op ultrasound, you may shower (**no tub-baths**) to wash off any stains. Do not scrub the wounds. You can rub the legs lightly with your fingertips. Use hibiclens, dial, or other bactericidal soap. Pat your legs dry when getting out of the shower, and then dry them completely with a hair-dryer on the "cool" setting. Leave the paper steri-strips on your wound (they will peel off in about 2 weeks). You may place a band-aid over the wounds if you wish.
- Starting postoperative day two or three (when the ace wraps are removed at the Vein Center), you should wear support stockings as much as you can for the next two weeks (daytime only). These can be purchased through our office for nearly one-half the usual cost.
- You may resume normal activities after 2 to 4 days, depending upon your leg discomfort. You will likely continue to experience some soreness along the course of the treated vein.
- You may resume a light exercise schedule within 2 weeks, again using common sense to adjust your schedule if your leg is still tender. If you exercise vigorously, walk several miles a day, or jog; please wear your support stockings for one month (especially during exercise).
- There will be some bruising, redness and tenderness in the areas that once contained varicose veins. Patients can often feel knots or bumps in the region the micro-phlebectomy or sclerotherapy has been performed (where the varicose veins had been). These knots are usually mildly tender, for they represent subcutaneous bruising and thrombosed varicose veins. This is a normal result of the procedure and is temporary. The bruising will not harm you, and is not the type of "blood clots" that can go to your lungs. They will slowly disappear as the veins dissolve away.
- If you travel by car or airplane for longer than one hour for the next six (6) months, please put on the support stockings while in transit. In addition, please walk around (in a parking lot or in the airplane aisle) for five minutes every 45 minutes to one hour. You do have an increased risk to have blood clots form in the deep veins of your leg if you are immobile for an extended period.
- You should avoid soaking in hot tubs or spas for at least 4 weeks after your operation. Swimming and tub baths are o.k. after 2 weeks.
- Resume your normal medications after your procedure. If you have questions about your medications, please ask Dr. Hayes.
- You will have a follow-up office visit and postoperative ultrasound within 1 week of your procedure. We will ask you to return for follow-up ultrasound at 6 weeks and 1 year from your procedure date in order to be sure that the long-term results are satisfactory to you and to Dr. Hayes. There is no charge to you for your follow-up visits.
- If you need additional spider vein treatments, Dr. Hayes will explain what can be done and how the spider veins can be eliminated. Because insurance companies consider spider veins a cosmetic problem, they will not cover this treatment. Dr. Hayes will give you the cost to treat each leg. He determines a one-time fee for each leg, which is good for at least three treatments up to one year. Dr. Hayes is known to be very fair regarding these treatments in order to provide an affordable service for you.
- You will be given a prescription for pain medication (Lortab or Darvocet) to be taken as needed. Be sure to take ibuprofen 600 mgs approximately every 8 hours for five days after the procedure to diminish postoperative inflammation (unless you are sensitive to ibuprofen). You will most likely find that your thigh is somewhat tender during the next few weeks. This is due to the inflammatory process that accompanies your body absorbing the ablated vein. This inflammatory process is quite normal. Ibuprofen taken as needed (not to exceed 400 mgs three times daily) will help relieve that tenderness.

*Most Frequently Asked Questions are answered in the **FAQ and Vein Myths** section of our website at www.veincenternorthtexas.com. If you have further questions, please feel free to call Dr. Hayes or his office staff at any time. You may also contact Dr. Hayes after office hours by using the Texoma Medical Center operator, or call him at his home. The contact numbers are listed below.*

Thank you,
Clint A. Hayes, M.D., F.A.C.S.

Vein Center of North Texas

102 Memorial Drive, Suite 101
Denison, Texas 75020

www.veincenternorthtexas.com

Vein Center of North Texas phone number **903-465-7070**
Texoma Medical Center telephone operator **903-416-4000**
Dr. Hayes' home number **903-465-2486**